



## Medical Necessity Guideline

<b>Medical Necessity Guideline (MNG) Title: Excision of Excessive Skin and Subcutaneous Tissue</b>		
<b>MNG #: 059</b>	<input checked="" type="checkbox"/> SCO <input checked="" type="checkbox"/> One Care	<b>Prior Authorization Needed?</b> <input checked="" type="checkbox"/> Yes <input type="checkbox"/> No
<b>Clinical:</b> <input checked="" type="checkbox"/>	<b>Operational:</b> <input type="checkbox"/>	<b>Informational:</b> <input type="checkbox"/>
<b>Medicare Benefit:</b> <input type="checkbox"/> Yes <input type="checkbox"/> No	<b>Approval Date:</b> 4/1/2021;	<b>Effective Date:</b> 06/19/2021
<b>Last Revised Date:</b>	<b>Next Annual Review Date:</b> 4/1/2022;	<b>Retire Date:</b>

**OVERVIEW:**

Rapid loss of massive amounts of weight results in excessive skin and subcutaneous tissue without potential for retraction. The excessive skin and subcutaneous tissue are most prevalent in the lower abdomen. In addition to cosmetic concerns, a large and heavy abdominal panniculus can interfere with normal activities of daily living, such as walking, climbing stairs, bathing or showering, and getting dressed. Rashes and skin irritation may occur on the opposing surfaces of the skin, particularly in warm weather. Occasionally secondary bacterial or fungal infections can complicate these skin rashes. Less commonly, folds of skin in other areas, such as the upper arms and thighs, may interfere with normal activities of daily living or cause rashes and skin irritations. Timing of surgery to remove excessive skin and subcutaneous tissue should be determined by the stabilization of the member’s weight. For members who have had bariatric surgery, this usually occurs 18 to 24 months after the procedure. Rarely, excess tissue may complicate wound healing due to traction or may need to be removed to expose other surgical areas or to minimize complications from a complex surgical procedure.

Commonwealth Care Alliance (CCA) considers approval for coverage of excision of excessive skin and subcutaneous tissue on an individual, case-by-case basis, in accordance with MassHealth guidelines and 130 CMR 409.000 and 130 CMR 450.204.

**Clinical Coverage Criteria:**

CCA bases its determination of medical necessity for excision of excessive skin and subcutaneous tissue on a combination of clinical data and the presence of indicators that would affect the relative risks and benefits of the procedure, including post-operative recovery. These criteria include all of the following:

1. The member has had massive weight loss. Massive weight loss varies based on individual clinical circumstances and is documented when:
  - a. the member has reached a body mass index (BMI) less than 30 kg/m<sup>2</sup> (from a pre-weight loss BMI ≥ 40 kg/m<sup>2</sup> or ≥ 35 kg/m<sup>2</sup> with at least one co-morbidity related to obesity, such as type 2 diabetes), or
  - b. the member has achieved at least a 100-pound weight loss, or
  - c. the member has achieved a 50% or greater excess body weight loss (%EBWL) or excess body mass index loss (%EBMIL). (See Appendix for formula for determining %EBWL and %EBMIL.)



## Medical Necessity Guideline

2. The member's weight has been stable for the preceding six months prior to the request. Stable weight is defined as less than 3% weight change.
  3. Standing photographs of the member clearly demonstrate excessive skin and subcutaneous tissue in the area being excised, and in the case of a request for panniculectomy, standing photographs (frontal and lateral) must clearly demonstrate that the panniculus covers the member's upper thigh crease (American Society of Plastic Surgeons (ASPS) Grade II).
  4. The excessive skin and subcutaneous tissue in the area being excised:
    - a. significantly interferes with the performance (impaired physical function) of normal activities of daily living (ADL), such as walking, climbing stairs, bathing or showering, and getting dressed, or
    - b. is causing recurrent (defined as two or more episodes over a 12-month period) skin or soft-tissue infections which have required medically supervised and documented antibiotic or antifungal therapy, which has not been effective.
- Note: Only in rare circumstances would excessive skin and subcutaneous tissue in the arms, thighs or buttocks, etc. cause significant impaired physical function or recurrent skin or soft tissue infections. Typically, these procedures are performed to improve appearance and are therefore cosmetic in nature.<sup>3</sup>
5. A comprehensive preoperative evaluation, including, but not limited to, obesity related comorbidities, such as diabetes and sleep apnea, and non-obesity related comorbidities, such as chronic obstructive pulmonary disease (COPD), nutritional status and psychosocial status has been conducted to identify the potential risks of the procedure.

Exception: In extraordinary circumstances panniculectomy may be performed to facilitate a complex surgical procedure such as hysterectomy and bilateral salpingoophorectomy performed via laparotomy. (The above criteria 1 and 2 related to weight loss do not apply in this case).

### LIMITATIONS/EXCLUSIONS:

(Documented limits or exclusions)

CCA does not consider excision of excessive skin and subcutaneous tissue to be medically necessary under certain circumstances. Examples of such circumstances include, but are not limited to, the following:

1. The member has difficulty in fitting clothes.
2. When a panniculectomy is being performed at the same time as bariatric surgery.
3. When a panniculectomy is being performed to prevent hernia occurrence or to prevent hernia recurrence in conjunction with a hernia repair, unless the member meets the criteria for panniculectomy stated in Section II. A. (1. – 5.) of these Guidelines.
4. When excision of excessive skin and subcutaneous tissue is being performed for cosmetic purposes, i.e., for the purpose of altering appearance, and is unrelated to physical disease or defect.



## Medical Necessity Guideline

### AUTHORIZATION:

#### Clinical Documentation Requirements

Requests for prior authorization for excision of excessive skin and subcutaneous tissue, including but not limited to panniculectomy (CPT code 15830), thighplasty (CPT 15832), and brachioplasty (CPT 15836), must be accompanied by clinical documentation that supports medical necessity. The quality of documentation is a critical factor in determination of medical necessity. In the absence of documentation supporting medical necessity, these procedures will be considered cosmetic.

- A. Documentation of medical necessity for each requested procedure must include the following (except for items 5 and 6 if the indication for tissue removal is not the result of massive weight loss):
1. The primary diagnosis name and current ICD-CM code pertinent to the clinical symptoms.
  2. The secondary diagnosis name and current ICD-CM code pertinent to comorbid condition(s).
  3. The member's comprehensive medical and surgical history, and when massive weight loss is the result of bariatric surgery, documentation must include immediate and late complications of the surgery, and post-surgical recovery.
  4. A list of the member's current prescribed and over-the-counter medications.
  5. Documentation of massive weight loss as defined in Section II.A.1. A BMI table (such as the one available on the National Institutes of Health (NIH) website: [https://www.nhlbi.nih.gov/health/educational/lose\\_wt/BMI/bmi\\_tbl.htm](https://www.nhlbi.nih.gov/health/educational/lose_wt/BMI/bmi_tbl.htm)) or a BMI calculator (such as the one available on the NIH website: <http://www.nhlbi.nih.gov/guidelines/obesity/BMI/bmicalc.htm>) can be used to determine the member's pre- and post-weight loss BMIs; see Appendix for formula for determining %EBWL and %EBMIL), and
  6. Documentation showing the member's weight has been stable for the preceding six months. Medical records documenting the member's weight over the preceding six months are required.
  7. Medical records documenting impaired physical function (if applicable).
8. Medical records documenting the assessment and treatment of 2 or more episodes of skin or soft-tissue infection over a 12-month period (if applicable).

### REGULATORY NOTES:

### RELATED REFERENCES:

MassHealth Guidelines for Medical Necessity Determination for Excision of Excessive Skin and Subcutaneous Tissue. Effective Date 12/22/2017. <https://www.mass.gov/doc/guidelines-for-medical-necessity-determination-for-excision-of-excessive-skin-and-subcutaneous/download>. Accessed 3/24/2021.

Local Coverage Determination (LCD) Cosmetic and Reconstructive Surgery (L34698).



## Medical Necessity Guideline

**ATTACHMENTS:**

<b>EXHIBIT A:</b>	
<b>EXHIBIT B</b>	

**REVISION LOG:**

REVISION DATE	DESCRIPTION

**APPROVALS:**

Douglas Hsu, MD

Vice President, Medical Policy and  
Utilization Management

\_\_\_\_\_  
**CCA Senior Clinical Lead [Print]**

\_\_\_\_\_  
**Title [Print]**

4/1/2021

\_\_\_\_\_  
**Signature**

\_\_\_\_\_  
**Date**

[Click here to enter text.](#)

\_\_\_\_\_  
**CCA Senior Operational Lead [Print]**

\_\_\_\_\_  
**Title [Print]**

\_\_\_\_\_  
**Signature**

\_\_\_\_\_  
**Date**

Lori Tishler, MD

Sr. Vice President, Medical Services

\_\_\_\_\_  
**CCA CMO or Designee [Print]**

\_\_\_\_\_  
**Title [Print]**

4/1/2021

\_\_\_\_\_  
**Signature**

\_\_\_\_\_  
**Date**