

UTILIZATION MANAGEMENT MEDICAL POLICY

POLICY: Oncology (Injectable – Programmed Death-Ligand 1) – Tecentriq Hybreza Utilization Management Medical Policy

- Tecentriq Hybreza™ (atezolizumab and hyaluronidase-tqjs subcutaneous injection – Genentech)

REVIEW DATE: 04/01/2026; selected revision 05/20/2026

OVERVIEW

Tecentriq Hybreza, a programmed death-ligand 1 (PD-L1) blocking antibody and hyaluronidase, is indicated for the treatment of the following:¹

- **Alveolar soft part sarcoma**, for the treatment of unresectable or metastatic disease in adults and pediatric patients ≥ 12 years of age who weigh ≥ 40 Kg.
- **Hepatocellular carcinoma**, in combination with bevacizumab, for the treatment of unresectable or metastatic hepatocellular carcinoma in adults who have not received prior systemic therapy.
- **Melanoma**, in combination with Cotellic® (cobimetinib tablets) and Zelboraf® (vemurafenib tablets), for the treatment of *BRAF V600* mutation-positive unresectable or metastatic disease as determined by an FDA-approved test in adults.
- **Muscle Invasive Bladder cancer**, as adjuvant treatment after cystectomy in adults who have circulating tumor DNA molecular residual disease (ctDNA MRD) as determined by an FDA-authorized test.
- **Non-small cell lung cancer (NSCLC), metastatic** disease in adults:
 - As a single agent, as adjuvant treatment following resection and platinum-based chemotherapy for patients with Stage II to IIIA disease whose tumors express PD-L1 on $\geq 1\%$ of tumor cells.
 - As a single-agent, for the first-line treatment of tumors with high PD-L1 expression (PD-L1 staining $\geq 50\%$ of tumor cells or PD-L1 staining of tumor infiltrating immune cells covering $\geq 10\%$ of the tumor area), with no anaplastic lymphoma kinase (*ALK*) or epidermal growth factor receptor (*EGFR*) genomic tumor aberrations.
 - In combination with bevacizumab, paclitaxel, and carboplatin, for the first-line treatment of metastatic non-squamous NSCLC with no *ALK* or *EGFR* genomic tumor aberrations.
 - In combination with paclitaxel protein-bound and carboplatin, for the first-line treatment of non-squamous metastatic NSCLC with no *ALK* or *EGFR* genomic tumor aberrations.
 - As a single-agent, for disease progression during or following platinum-containing chemotherapy. Patients with *EGFR* or *ALK* genomic tumor aberrations should have disease progression on FDA-approved therapy for these aberrations prior to receiving Tecentriq Hybreza.
- **Small cell lung cancer**, in adults:
 - In combination with carboplatin and etoposide, for the first-line treatment of extensive-stage disease.
 - In combination with Zepzelca® (lurbinectedin intravenous infusion) for maintenance treatment of extensive-stage disease that has not progressed after first-line induction therapy with Tecentriq Hybreza or Tecentriq® (atezolizumab intravenous infusion), carboplatin and etoposide.

Guidelines

The use of Tecentriq is recommended across multiple National Comprehensive Cancer Network (NCCN) guidelines and the NCCN Compendium.²⁻¹⁵ Recommendations with a category of evidence of 2B or higher support the approval criteria outlined below.

POLICY STATEMENT

Prior Authorization is recommended for medical benefit coverage of Tecentriq Hybreza. Approval is recommended for those who meet the **Criteria** and **Dosing** for the listed indications. Extended approvals are allowed if the patient continues to meet the Criteria and Dosing. Requests for doses outside of the established dosing documented in this policy will be considered on a case-by-case basis by a clinician (i.e., Medical Director or Pharmacist). All approvals are provided for the duration noted below. Because of the specialized skills required for evaluation and diagnosis of patients treated with Tecentriq Hybreza as well as the monitoring required for adverse events and long-term efficacy, approval requires Tecentriq Hybreza to be prescribed by or in consultation with a physician who specializes in the condition being treated.

Automation: None.

RECOMMENDED AUTHORIZATION CRITERIA

Coverage of Tecentriq Hybreza is recommended in those who meet one of the following criteria:

FDA-Approved Indications

-
1. **Alveolar Soft Part Sarcoma.** Approve for 1 year if the patient meets BOTH of the following (A, B, and C):
- A) Patient is ≥ 12 years of age; AND
 - B) Patient weighs ≥ 40 Kg; AND
 - C) The medication is prescribed by or in consultation with an oncologist.

Dosing. Approve 1,875 mg of atezolizumab and 30,000 units of hyaluronidase (15 mL) administered subcutaneously no more frequently than once every 3 weeks.

-
2. **Hepatocellular Carcinoma.** Approve for 1 year if the patient meets ALL of the following (A, B, C, and D):
- A) Patient is ≥ 18 years of age; AND
 - B) Patient meets ONE of the following (i or ii):
 - i. Patient meets BOTH of the following (a and b):
 - a) The medication is used for first-line therapy; AND
 - b) According to the prescriber, the patient has ONE of the following [(1) or (2)]:
 - (1) Liver confined, unresectable disease and is deemed ineligible for transplant; OR
 - (2) Extrahepatic/metastatic disease and is deemed ineligible for resection, transplant, or locoregional therapy; OR
 - ii. The medication will be used as subsequent-line therapy; AND
 - C) The medication will be used in combination with bevacizumab; AND
 - D) The medication is prescribed by or in consultation with an oncologist.
-

Dosing. Approve 1,875 mg of atezolizumab and 30,000 units of hyaluronidase (15 mL) administered subcutaneously no more frequently than once every 3 weeks.

-
3. **Melanoma.** Approve for 1 year if the patient meets ALL of the following (A, B, C, and D):
- A) Patient is ≥ 18 years of age; AND
 - B) Patient has *BRAF V600* mutation-positive disease; AND
 - C) The medication will be used in combination with Cotellic (cobimetinib tablets) and Zelboraf (vemurafenib tablets); AND
 - D) The medication is prescribed by or in consultation with an oncologist.

Dosing. Approve 1,875 mg of atezolizumab and 30,000 units of hyaluronidase (15 mL) administered subcutaneously no more frequently than once every 3 weeks.

-
4. **Muscle Invasive Bladder Cancer:** Approve for 1 year if the patient meets ALL of the following (A, B, C, D, and E):
- A) Patient is ≥ 18 years of age; AND
 - B) Patient had a cystectomy; AND
 - C) Patient has circulating tumor DNA molecule residual disease (ctDNA MDR); AND
 - D) The medication is used as adjuvant therapy; AND
 - E) The medication is prescribed by or in consultation with an oncologist.

Dosing. Approve 1,875 mg of atezolizumab and 30,000 units of hyaluronidase (15 mL) administered subcutaneously no more frequently than once every 3 weeks.

-
5. **Non-Small Cell Lung Cancer – Adjuvant.** Approve for 1 year if the patient meets ALL of the following (A, B, C, and D):
- A) Patient is ≥ 18 years of age; AND
 - B) The tumor is negative for the following actionable biomarkers: epidermal growth factor receptor (*EGFR*) exon 19 deletion or exon 21 L858R, anaplastic lymphoma kinase (*ALK*), *RET*, or *ROS1*; AND
 - C) Patient meets ALL of the following (i, ii, and iii):
 - i. Patient has completely resected or node positive disease; AND
 - ii. The tumor expresses programmed death-ligand 1 (PD-L1) $\geq 1\%$; AND
 - iii. Patient has received previous adjuvant chemotherapy; AND
 - D) The medication is prescribed by or in consultation with an oncologist.

Dosing. Approve 1,875 mg of atezolizumab and 30,000 units of hyaluronidase (15 mL) administered subcutaneously no more frequently than once every 3 weeks.

-
6. **Non-Small Cell Lung Cancer – Recurrent, Advanced, or Metastatic.** Approve for 1 year if the patient meets ALL of the following (A, B, and C):
- A) Patient is ≥ 18 years of age; AND
 - B) Patient meets ONE of the following (i or ii):
 - i. The tumor is positive for epidermal growth factor receptor (*EGFR*) exon 19 deletion or exon 21 L858R, anaplastic lymphoma kinase (*ALK*), *RET*, or *ROS1*; OR

- ii. The tumor is negative for the following actionable biomarkers: epidermal growth factor receptor (*EGFR*) exon 19 deletion or exon 21 L858R, anaplastic lymphoma kinase (*ALK*), *RET*, or *ROS1* and meets ONE of the following (a, b, c, or d):

- a) The medication is used as first-line therapy and the tumor is positive for ONE of the following[(1), (2), or (3)]:
(1) *EGFR* exon 20 mutation; OR
(2) *NRG1* gene fusion positive; OR
(3) *ERBB2* (*HER2*) mutation positive; OR
- b) The medication is used for first-line or subsequent therapy and the tumor is positive for ONE of the following [(1), (2), or (3)]:
(1) *BRAF V600E* mutation; OR
(2) *NTRK1/2/3* gene fusion; OR
(3) *MET* exon 14 skipping mutation positive; OR
- c) The medication is used as subsequent therapy and the tumor is *EGFR S768I*, *L861Q*, and/or *G719X* mutation positive; OR
- d) The medication is used as first-line or continuation maintenance therapy and the tumor has no actionable mutations; AND

Note: The tumor does NOT have the following mutations: *EGFR* exon 19 deletion, *EGFR* exon 21 L857R, *EGFR* S768I, *EGFR* L861Q, *EGFR* G719X, *EGFR* exon 20 insertion, *ALK* rearrangement, *ROS1* rearrangement, *BRAF* V600E, *NTRK* 1/2/3 gene fusion, *MET*ex14 skipping, *RET* rearrangement, *ERBB2* (*HER2*), and *NRG1* gene fusion.

- C) The medication is prescribed by or in consultation with an oncologist.

Dosing. Approve 1,875 mg of atezolizumab and 30,000 units of hyaluronidase (15 mL) administered subcutaneously no more frequently than once every 3 weeks.

7. Small Cell Lung Cancer. Approve for 1 year if the patient meets BOTH of the following (A and B):

- A) Patient is ≥ 18 years of age; AND
B) The medication is prescribed by or in consultation with an oncologist.

Dosing. Approve 1,875 mg of atezolizumab and 30,000 units of hyaluronidase (15 mL) administered subcutaneously no more frequently than once every 3 weeks.

Other Uses with Supportive Evidence

8. Cervical Cancer. Approve for 1 year if the patient meets ALL of the following (A, B, and C):

- A) Patient is ≥ 18 years of age; AND
B) Patient has persistent, recurrent, or metastatic disease; AND
C) The medication is prescribed by or in consultation with an oncologist.

Dosing. Approve 1,875 mg of atezolizumab and 30,000 units of hyaluronidase (15 mL) administered subcutaneously no more frequently than once every 3 weeks.

9. Chronic Lymphocytic Leukemia/Small Lymphocytic Lymphoma. Approve for 1 year if the patient meets ALL of the following (A, B, C, and D):

- A) Patient is ≥ 18 years of age; AND

- B) Patient has histologic Richter transformation; AND
- C) The medication is used in combination with Venetoclax and Gazyva (obinutuzumab intravenous injection); AND
- D) The medication is prescribed by or in consultation with an oncologist.

Dosing. Approve 1,875 mg of atezolizumab and 30,000 units of hyaluronidase (15 mL) administered subcutaneously no more frequently than once every 3 weeks.

10. Colon and Rectal Cancer: Approve for 1 year if the patient meets ALL of the following (A, B, and C)

- A) The patient is ≥ 18 years of age; AND
- B) Patient meets ONE of the following (i or ii):
 - i. Patient has microsatellite instability-high (MSI-H) or mismatch repair deficient (dMMR); OR
 - ii. Patient has polymerase epsilon/delta (POLE/POLD1) mutation positive with ultra-hypermutated phenotype; AND

Note: Ultra-hypermutated phenotype is defined as tumor mutational burden > 50 mutations/megabase.
- C) The medication is prescribed by or in consultation with an oncologist.

Dosing. Approve 1,875 mg of atezolizumab and 30,000 units of hyaluronidase (15 mL) administered subcutaneously no more frequently than once every 3 weeks.

11. Mesothelioma. Approve for 1 year if the patient meets ALL of the following (A, B, C, D, and E):

- A) Patient is ≥ 18 years of age; AND
- B) The medication is used as subsequent therapy; AND
- C) The medication is used in combination with bevacizumab; AND
- D) Patient has ONE of the following (i, ii, or iii):
 - i. Malignant peritoneal mesothelioma; OR
 - ii. Pericardial mesothelioma; OR
 - iii. Tunica vaginalis testis mesothelioma; AND
- E) The medication is prescribed by or in consultation with an oncologist.

Dosing. Approve 1,875 mg of atezolizumab and 30,000 units of hyaluronidase (15 mL) administered subcutaneously no more frequently than once every 3 weeks.

12. Small Bowel Adenocarcinoma. Approve for 1 year if the patient meets ALL of the following (A, B, C, and D):

- A) Patient is ≥ 18 years of age; AND
- B) Patient meets ONE of the following (i or ii):
 - i. Patient has advanced or metastatic disease; OR
 - ii. The medication is used as neoadjuvant or adjuvant therapy; AND
- C) Patient meets ONE of the following (i or ii):
 - i. Patient has deficient mismatch repair/microsatellite instability-high (dMMR/MSI-H) disease; OR
 - ii. The disease is polymerase epsilon/delta (POLE/POLD1) mutation positive with ultra-hypermutated phenotype; AND

Note: Ultra-hypermutated phenotype is defined as tumor mutational burden > 50 mutations/megabase.

D) The medication is prescribed by or in consultation with an oncologist.

Dosing. Approve 1,875 mg of atezolizumab and 30,000 units of hyaluronidase (15 mL) administered subcutaneously no more frequently than once every 3 weeks.

13. Thymic Carcinoma. Approve for 1 year if the patient meets BOTH of the following (A, B, and C):

A) Patient is ≥ 18 years of age; AND

B) The medication is used in combination with carboplatin and paclitaxel; AND

C) The medication is prescribed by or in consultation with an oncologist.

Dosing. Approve 1,875 mg of atezolizumab and 30,000 units of hyaluronidase (15 mL) administered subcutaneously no more frequently than once every 3 weeks.

14. Urothelial Carcinoma. Approve for 1 year if the patient meets ALL of the following (A, B, and C):

A) Patient is ≥ 18 years of age; AND

B) Patient meets ONE of the following (i or ii):

i. The tumor expresses programmed death-ligand 1 (PD-L1); OR

ii. Patient is ineligible for any platinum-containing chemotherapy regardless of PD-L1 expression; AND

C) The medication is prescribed by or in consultation with an oncologist.

Dosing. Approve 1,875 mg of atezolizumab and 30,000 units of hyaluronidase (15 mL) administered subcutaneously no more frequently than once every 3 weeks.

CONDITIONS NOT RECOMMENDED FOR APPROVAL

Coverage of Tecentriq Hybreza is not recommended in the following situations:

1. Coverage is not recommended for circumstances not listed in the Recommended Authorization Criteria. Criteria will be updated as new published data are available.

REFERENCES

1. Tecentriq Hybreza subcutaneous injection [prescribing information]. South San Francisco, CA: Genentech; May 2026.
2. The NCCN Drugs & Biologics Compendium. © 2026 National Comprehensive Cancer Network. Available at: <http://www.nccn.org>. Accessed on March 19, 2026. Search term: atezolizumab and hyaluronidase.
3. The NCCN Soft Tissue Sarcoma Clinical Practice Guidelines in Oncology (version 3.2026 – March 12, 2026). © 2026 National Comprehensive Cancer Network. Available at: <http://www.nccn.org>. Accessed March 24, 2026.
4. The NCCN Non-Small Cell Lung Cancer Clinical Practice Guidelines in Oncology (version 5.2026 – March 13, 2026). © 2026 National Comprehensive Cancer Network. Available at: <http://www.nccn.org>. Accessed on March 20, 2026.
5. The NCCN Bladder Cancer Clinical Practice Guidelines in Oncology (version 1.2026 – March 16, 2026). © 2026 National Comprehensive Cancer Network. Available at: <http://www.nccn.org>. Accessed on March 24, 2026.
6. The NCCN Small Cell Lung Cancer Clinical Practice Guidelines in Oncology (version 2.2026 – September 16, 2025). © 2025 National Comprehensive Cancer Network. Available at: <http://www.nccn.org>. Accessed on March 19, 2026.
7. The NCCN Hepatocellular Carcinoma Clinical Practice Guidelines in Oncology (version 1.20265 – March 10, 2026). © 2026 National Comprehensive Cancer Network. Available at: <http://www.nccn.org>. Accessed on March 20, 2026.
8. The NCCN Melanoma: Cutaneous Clinical Practice Guidelines in Oncology (version 1.2026 – February 17, 2026). © 2026 National Comprehensive Cancer Network. Available at: www.nccn.org. Accessed on March 20, 2026.

9. The NCCN Mesothelioma: Peritoneal Clinical Practice Guidelines in Oncology (version 2.2026 – October 3, 2025). © 2025 National Comprehensive Cancer Network. Available at: <http://www.nccn.org>. Accessed March 19, 2026.
10. The NCCN Cervical Cancer Clinical Practice Guidelines in Oncology (version 2.2026 – November 10, 2025). © 2025 National Comprehensive Cancer Network. Available at: <http://www.nccn.org>. Accessed March 19, 2026.
11. The NCCN Colon Cancer Clinical Practice Guidelines in Oncology (version 4.2025 – June 27, 2025). © 2025 National Comprehensive Cancer Network. Available at: <http://www.nccn.org>. Accessed August 14, 2025.
12. The NCCN Thymomas and Thymic Carcinoma Clinical Practice Guidelines in Oncology (version 1.2026 – October 3, 2025). © 2025 National Comprehensive Cancer Network. Available at: <http://www.nccn.org>. Accessed March 19, 2026.
13. The NCCN Chronic Lymphocytic Leukemia/Small Lymphocytic Lymphoma Clinical Practice Guidelines in Oncology (version 2.2026 – December 22, 2025). © 2025 National Comprehensive Cancer Network. Available at: <http://www.nccn.org>. Accessed March 20, 2026.
14. The NCCN Small Bowel Adenocarcinoma Clinical Practice Guidelines in Oncology (version 1.2026 – January 22, 2026). © 2026 National Comprehensive Cancer Network. Available at: <http://www.nccn.org>. Accessed March 24, 2026.
15. The NCCN Rectal Cancer Clinical Practice Guidelines in Oncology (version 1.2026 – March 5, 2026). © 2026 National Comprehensive Cancer Network. Available at: <http://www.nccn.org>. Accessed March 24, 2026.

HISTORY

Type of Revision	Summary of Changes	Review Date
New Policy	--	09/25/2024
Early Annual Revision	Hepatocellular Carcinoma: Duration of approval was changed from 1 year to approve for the duration noted. Requirements that the patient has Child-Pugh Class A or B liver function and patient has not received prior systemic therapy were removed. Option for approval that the patient has unresectable or metastatic hepatocellular carcinoma and is not a surgical candidate; and patient has liver confined disease, inoperable by performance status, comorbidity, or with minimal or uncertain extrahepatic disease were removed. Added patient has undergone resection or ablation therapy, patient is at high-risk of recurrence, and medication is used for adjuvant therapy as new option for approval with duration of up to 1 year (total). Added medication is used first-line and patient has liver confined, unresectable disease and is deemed ineligible for transplant; or extrahepatic/metastatic disease and are deemed ineligible for resection, transplant, or locoregional therapy, with a 1 year duration of approval. Non-Small Cell Lung Cancer: Removed <i>KRAS</i> from list of actionable mutations and added may be <i>KRAS G12C</i> mutation positive to Note. Removed patient's tumor expresses programmed death-ligand 1 (PD-L1) $\geq 1\%$ and medication will be used in combination with chemotherapy as options for approval. Added descriptor first-line to the medication will be used as first-line or continuation maintenance therapy. Added requirement that the medication is used as first-line or continuation maintenance therapy. Added <i>EGFR</i> exon 19 deletion, exon 21 <i>L858R</i> mutation, <i>ALK</i> rearrangement, <i>RET</i> rearrangement, and <i>ROS1</i> rearrangement negative to requirement that patient has recurrent, advanced, or metastatic non-squamous cell NSCLC, is <i>EGFR</i> exon 19 deletion, exon 21 <i>L858R</i> mutation, <i>ALK</i> rearrangement, <i>RET</i> rearrangement, and <i>ROS1</i> rearrangement negative and meets ONE of the following. Removed <i>KRAS G12C</i> mutation and added <i>NRG1</i> gene fusion to requirement that the tumor is <i>EGFR</i> exon 20 mutation positive, <i>NRG1</i> gene fusion positive or <i>ERBB2 (HER2)</i> mutation positive. Removed <i>RET</i> rearrangement positive from requirement that the tumor is <i>BRAF V600E</i> mutation positive, <i>NTRK1/2/3</i> gene fusion positive, or <i>MET</i> exon 14 skipping mutation positive. Removed <i>EGFR</i> exon 19 deletion, exon 21 <i>L858R</i> positive, <i>ALK</i> rearrangement positive, or <i>ROS1</i> rearrangement positive from requirement that the tumor is <i>EGFR S768I</i>, <i>L861Q</i>, and/or <i>G719X</i> mutation positive. Removed examples of targeted drug therapy. Added option for approval for patients with squamous cell NSCLC, performance status of 3, and medication is used as a single agent. Added requirements for adjuvant therapy: patient has completed resected disease and patients is negative for <i>EGFR</i> exon 19 deletion, exon 21 <i>L858R</i> mutations and <i>ALK</i> rearrangements. Cervical Cancer: Requirement that the patient has small neuroendocrine carcinoma of the cervix was removed. Urothelial Carcinoma: Added new condition of approval.	01/22/2025

HISTORY (CONTINUED)

Type of Revision	Summary of Changes	Review Date
Early Annual Revision	Alveolar Soft Part Sarcoma: The requirements the patient has unresectable or metastatic disease and the medication is used as a single agent were removed. Hepatocellular Carcinoma: The approval duration was modified to approve for 1 year for all approval options. The options for approval that the patient has undergone resection or ablation therapy, has high-risk of recurrence, and the medication is used as adjuvant therapy has been removed. The medication will be used as subsequent-line therapy was added as an approval option. Melanoma: The requirements that the patient has unresectable or metastatic melanoma and that the medication will be used as subsequent therapy have been removed. Non-Small Cell Lung Cancer – Adjuvant: The condition of approval was changed to as listed. Previously, all non-small cell lung cancer (NSCLC) was addressed more generally under NSCLC. The approval duration was modified to approve for 1 year. A requirement was added that the tumor is negative for the following actionable biomarkers: epidermal growth factor receptor (EGFR) exon 19 deletion or exon 21 L858R, anaplastic lymphoma kinase (ALK), RET, and ROS1. Non-Small Cell Lung Cancer – Recurrent, Advanced, or Metastatic Disease: The indication was changed to as listed. Previously, all non-small cell lung cancer (NSCLC) was addressed more generally under NSCLC. The approval duration was modified to approve for 1 year for all approval options. A requirement was added that the tumor is positive for epidermal growth factor receptor (EGFR) exon 19 deletion or exon 21 L858R, anaplastic lymphoma kinase (ALK), RET, or ROS1. The differentiation of the different types of NSCLC (non-squamous cell NSCLC and squamous cell NSCLC) were removed as options for approval. For all lines of therapy, removed the medication will be used in combination with chemotherapy as an option for approval. For subsequent therapy, the patient has received targeted drug therapy for the specific mutation and that the patient has not progressed on programmed death receptor-1 (PD-1) or programmed death-ligand 1 (PD-L1) were removed as options of approval. The patient has squamous cell NSCLC, is performance status 3, and the medication is used as a single-agent have been removed as approval options. The patient has completely resected disease and the tumor expresses programmed death-ligand 1 (PD-L1) $\geq 1\%$ as determined by an approved test were removed as approval options. For first-line therapy or as continuation maintenance therapy, a requirement was added that the tumor has no actionable mutations; a Note was added listing the following actionable mutations: EFGR exon 19 deletion, EFGR exon 21 L858R, EFGR S768I, EGFR L861Q, EGFR G719X, EGFR exon 20 insertion, ALK rearrangement, ROS1 rearrangement, BRAF V600E, NTRK 1/2/3 gene fusion, METex14 skipping, RET rearrangement, ERBB2 (HER2), and NRG1 gene fusion. For first-line and continuation maintenance therapy; the patient's tumor expresses programmed death-ligand 1 (PD-L1) $\geq 50\%$ as determined by an approved test was removed as an approval option. Colon Cancer: This was added as a new condition of approval. Urothelial Carcinoma: The options that the patient is currently receiving Tecentriq for the treatment of urothelial carcinoma and according to the prescriber, the patient is deriving benefit from Tecentriq were removed.	09/03/2025
Update	Overview: Updated the overview to reflect changes in the Small-Cell Lung Cancer indication. No criteria changes.	10/08/2025
Selected Revision	Alveolar Soft Part Sarcoma: The requirement that the patient is ≥ 18 years of age was modified to the patient is ≥ 12 years of age. The requirement that the patient weighs ≥ 40 Kg was added.	12/17/2025

HISTORY (CONTINUED)

Type of Revision	Summary of Changes	Review Date
Early Annual Revision	Cervical Cancer: The requirement that the medication is used in combination with chemotherapy and the corresponding Note was removed. Chronic Lymphocytic Leukemia/Small Lymphocytic Lymphoma: This was added as a new condition of approval. Colon and Rectal Cancer.: The condition of approval was changed to as listed. Previously listed as Colon Cancer. The definition of ultra-hypermutated phenotype was moved to a Note. Small Bowel Adenocarcinoma: This was added as a new condition of approval. Thymic Carcinoma: This was added as a new condition of approval. Urothelial Carcinoma: The requirement that the medication is used as first-line therapy was removed. The option of approval that the patient is ineligible for cisplatin and tumor expresses programmed death-ligand 1 (PD-L1) tumor infiltrating immune cells covering $\geq 5\%$ of tumor area was modified to the tumor expresses programmed death-ligand 1 (PD-L1).	04/01/2026
Selected Revision	Muscle Invasive Bladder Cancer: This was added as a new condition of approval.	05/20/2026