

Esketamine Authorization Request

Request Information

Expedited Request (by checking this box I certify that this request meets the below criteria for being Expedited and I will supply justification*)

Criteria for Expedited: Waiting for a decision within the standard time frame (up to 14 calendar days) could place the member's life, health, or ability to regain maximum function in serious jeopardy

***Justification for Expedited:**
(Attach pages if add'tl space is needed)

Member Name:		DOB:	Policy #
Facility Name:		Facility TIN:	
MD Performing Treatment:	Fax #	<input type="checkbox"/> Out of Network	
Provider Contact Name:		Contact Phone:	
Requested Procedure Code G2082 = G2083 =		Primary Diagnosis (Diagnosis code):	
Service Start Date:	Service End Date:	Frequency:	

Please select ONE:

- Initial authorization
- Authorization renewal (*skip to section on page 2 titled "For Authorization Renewals Only"*)

Please select ONE:

- Major Depressive Disorder and acute suicidal ideation or behavior confirmed by a psychiatrist (urgent)
- Major Depressive Disorder with treatment resistant depression confirmed by psychiatrist **OR**
- None of the above please specify:

Please select YES or NO:

- Previous treatment with more than 2 antidepressants for at least 6 weeks each? Yes No
- Continued depression after treatment? Yes No
- Pre-Esketamine treatment depression rating scale: GDS _____, PHQ-9 _____, BDI _____, HAM-D, MADRS _____, QIDS _____, or IDS-SR
- Is the drug being prescribed Esketamine? Yes No If no, indicate the drug to be prescribed:
- Esketamine treatment to be used in combination with an oral antidepressant? Yes No
- Is treatment to be provided in a hospital setting by a provider? Yes No
- If no, describe setting
- Is there a history of psychosis? Yes No
- If yes, does the prescriber believe the benefits of Esketamine outweigh the risks? Yes No

Please select **ALL** that apply from the below indicators for Esketamine treatment:

- No history of aneurysmal vascular disease, arteriovenous malformation, or intracranial hemorrhage
- Monitoring planned after each administration
- Risks of sedation and dissociation after administration discussed with patient or caregiver
- Risk of abuse and misuse discussed with patient or caregiver
- Risk of increased suicidal thoughts and behavior discussed with patient or caregiver
- Not currently pregnant or breastfeeding and risks of pregnancy/breastfeeding discussed with patient or caregiver, or pregnancy testing not indicated
- Does not have a hypersensitivity to Esketamine or any excipients
- Does not have a current substance use disorder, unless in remission
- Has **NOT** had previous treatment that was determined not to reduce symptoms or be efficacious

OR

- Other clinical information

Complete the Following for Authorization Renewals Only

Please select **ALL** that apply from the below indicators for Continuation of Esketamine treatment:

- Drug being prescribed is Esketamine If no, indicate the drug to be prescribed:
- Administration and monitoring of Esketamine is to be provided in a hospital setting by a provider
If no, describe setting
- Risk of abuse and misuse discussed with patient or caregiver
- Condition improved with treatment
Depression scale (initial & most recent): GDS , PHQ-9 , BDI , HAM-D , MADRS
QIDS , or IDS-SR
- Manageable or no side effects
- Treatment used in combination with oral depressant
- Does not have a current substance use disorder, unless in remission

OR

Other clinical information:

Additional Comments: